

## Cosmology

## Part A (problems to be handed in)

1. If you have an error of 0.1 magnitudes in the distance modulus,  $m-M$ , what will be the error in the distance,  $D$ . (7 points)

2. The surface brightness of an astronomical object is defined as its observed flux divided by its observed angular scale, i.e.  $\Sigma \sim F_{\text{obs}}/(\theta_{\text{obs}})^2$ . What is the redshift evolution of  $\Sigma(z)$  for a class of objects that are both standard candle and ruler? (7 points)

3. Plot the redshift evolution of the mass of various “ $\nu\sigma$  halos”, i.e.  $M_{\nu\sigma}(z)$ , where  $\nu$  is defined via

$$\nu = \frac{\delta_c}{D(z) \sigma(M_{\nu\sigma})}$$

Here  $D(a=1/(1+z))$  is the growth factor that can be approximated as

$$D(a) = \left[ \frac{5a}{2} \Omega_m(a) \left[ \Omega_m^{4/7}(a) - \Omega_\Lambda(a) + \left( 1 + \frac{\Omega_m(a)}{2} \right) \left( 1 + \frac{\Omega_\Lambda(a)}{70} \right) \right] \right]^{-1}$$

and

$$\sigma^2(M) = \frac{1}{2\pi^2} \int_0^\infty P_0(k) \hat{W}^2(kR) k^2 dk$$

where  $M$  and  $R$  are related in a way as given in the lecture notes. Please use a scale-free cosmology characterized by  $P_0(k) = A k^n$ ,  $\Omega_{m,0} = 1.0$ , and  $\Omega_{\Lambda,0} = 0$ . Vary both  $\nu$  and  $n$  in a reasonable range.

**hints:** During the calculation it makes sense to define a ‘typical collapsing mass’  $M_* = \sigma_0^{n+3} \frac{4\pi \langle \rho \rangle}{3}$

which you can safely assume to be  $M_* = 10^{13} M_\odot$ , irrespective of  $n$ .

Note that  $\sigma_0$  captures everything that is constant, even though still depends on  $n$  (but see previous hint!)

As explained in class, for each redshift  $z$  you need to find  $M_{\nu\sigma}(z)$  for which it helps to first find an

analytical expression for  $\sigma^2(M) = \frac{1}{2\pi^2} \int_0^\infty P_0(k) \hat{W}^2(kR) k^2 dk$  (8 points)

4. You are given some preliminary data from a GW observation:

i) The spectral noise density of the detector is approximately  $S_n(f) \sim 0.00938426 * (f/1\text{Hz})^{-20}$  sec,

ii) The maximum frequency of observations is  $f_{\text{max}} \sim 200\text{Hz}$ ,

iii) The signal-to-noise ratio of the observation was  $S/N \sim 8$ ,

iv) The frequencies of the GW for the last 0.1secs of the observation before the merger were (time in secs, freq. in Hz):

{0.1, 81.8089}, {0.09, 85.1059}, {0.08, 88.9492}, {0.07, 93.5166}, {0.06, 99.0818}, {0.05, 106.093}, {0.04, 115.353}, {0.03, 128.493}, {0.02, 149.594}, {0.01, 193.999}

Then:

1. Assuming circular orbits (averaged over the inclination), find the chirp mass and distance (in Mpc) of the system.

2. Finally, assuming the inclination angle was  $\pi/3$ , plot the reconstructed strains as a function of time for the last 0.1secs before the merger. (8 points)

## Cosmology

## Part B (problems to be discussed in class)

- 1) Explain the Saha equation
- 2) Derive the relation between (adiabatic) matter and temperature perturbations
- 3) Derive the scaling relations:

radiation domination:  $\delta_m \sim \ln(a)$

matter domination:  $\delta_m \sim a$

$\Lambda$  domination:  $\delta_m \sim 1/a^2$

- 4) Explain the (idea of) the Press-Schechter mass function
- 5) Explain the dependence of the CMB peaks on the parameters  $\Omega_m, \Omega_b, \Omega_k$
- 6) Calculate the sound horizon at recombination for  $\Omega_m=0.3$  and  $\Omega_m=1$ . Discuss what you found.
- 7) Explain the behavior of the matter power spectrum  $P(k)$